

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) (rEGP40/1372), 0.2mg/mL, IgG1, Clone: [rEGP40/1372], Mouse, Monoclonal BOT-BNUB2172-500

Article Name	Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) (rEGP40/1372), 0.2mg/mL, IgG1, Clone: [rEGP40/1372], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB2172-500
Supplier Catalog Number	BNUB2172-500
Alternative Catalog Number	BOT-BNUB2172-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Recombinant human EpCAM protein fragment from extracellular domain (around aa77-202) (exact sequence is proprietary)
Product Description	EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[rEGP40/1372]

Molecular Weight	40-43 kDa
Isotype	IgG1
UniProt	P16422
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user