

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) (rBCL2/782), 0.2mg/mL, Clone: [rBCL2/782], Mouse, Monoclonal BOT-BNUB2221-500

Article Name	Bcl-2 (Apoptosis & Follicular Lymphoma Marker) (rBCL2/782), 0.2mg/mL, Clone: [rBCL2/782], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB2221-500
Supplier Catalog Number	BNUB2221-500
Alternative Catalog Number	BOT-BNUB2221-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Recombinant full-length human BCL2 protein
Product Description	This antibody recognizes a protein of 25-26 kDa, identified as Bcl-2. This antibody shows no cross-reaction with Bcl-x or Bax proteins. Bcl-2 suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural ...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[rBCL2/782]
Molecular Weight	25-26 kDa

UniProt	P10415
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 1 mM EDTA Buffer pH 8.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user