

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

B7-H4 (Immuno-Inhibitory Protein) (B7H4/2652R), 0.2mg/mL, Clone: [B7H4/2652R], Rabbit, Monoclonal BOT-BNUB2652-100

Article Name	B7-H4 (Immuno-Inhibitory Protein) (B7H4/2652R), 0.2mg/mL, Clone: [B7H4/2652R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB2652-100
Supplier Catalog Number	BNUB2652-100
Alternative Catalog Number	BOT-BNUB2652-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	FC, IF, IHC
Species Reactivity	Human
Immunogen	Recombinant human B7-H4 protein fragment (exact sequence is proprietary)
Product Description	B7-H4 (B7-homolog 4), belongs to the B7 immunoglobulin superfamily of ligand-lymphocyte interacting proteins. Expressed primarily on the membrane of lymphoid cells, B7-H4 is an immuno-inhibitory protein that interacts with receptors on the surface of...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[B7H4/2652R]
Molecular Weight	35 kDa

UniProt	Q7Z7D3
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user