

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mucin 5AC (MUC5AC)(45M1), 1mg/mL, Clone: [45M1], Mouse, Monoclonal BOT-BNUM0031-50

Article Name	Mucin 5AC (MUC5AC)(45M1), 1mg/mL, Clone: [45M1], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM0031-50
Supplier Catalog Number	BNUM0031-50
Alternative Catalog Number	BOT-BNUM0031-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	ELISA, FC, IHC, WB
Species Reactivity	Feline, Gallus, Human, Mammal, Monkey, Mouse, Porcine, Rabbit, Rat
Immunogen	M1 mucin preparation from the fluid of an ovarian mucinous cyst belonging to an O Le(a-b) patient
Product Description	This MAb recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the C-terminal cysteine rich part of the peptide core of MUC5AC. The epitope is destroyed by beta-mercaptoethanol but not by period...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[45M1]
Molecular Weight	>1,000 kDa

UniProt	P98088
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Flow cytometry: 0.5-1 ug/million cells in 0.1 mL Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 1-2 ug/mL Does not react with cow, others not known Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user