

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Granulocyte-Colony Stimulating Factor (G-CSF) (CSF3)(CSF3/900), CF740 conjugate, 0.1mg/mL, IgG1, Clone: [CSF3/900], CF 740, Mouse, Monoclonal BOT-BNC740900-500

| | |
|----------------------------|---|
| Article Name | Granulocyte-Colony Stimulating Factor (G-CSF) (CSF3)(CSF3/900), CF740 conjugate, 0.1mg/mL, IgG1, Clone: [CSF3/900], CF 740, Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC740900-500 |
| Supplier Catalog Number | BNC740900-500 |
| Alternative Catalog Number | BOT-BNC740900-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human, Primate |
| Immunogen | Human recombinant full-length CSF3 protein |
| Conjugation | CF 740 |
| Product Description | This MAb recognizes granulocyte-colony stimulating factor (G-CSF) in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell types. Markers of myeloid cells are useful in the identification of different levels of cellular dif... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [CSF3/900] |

| | |
|-------------------|--|
| Molecular Weight | 19 kDa |
| Isotype | IgG1 |
| UniProt | P09919 |
| Buffer | PBS, 0.1% rBSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |