

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Lactoferrin protein, His tag, Unconjugated GTX00078-PRO

Article Name	Goat Lactoferrin protein, His tag, Unconjugated
Biozol Catalog Number	GTX00078-PRO
Supplier Catalog Number	GTX00078-pro
Alternative Catalog Number	GTX00078-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Goat
Conjugation	Unconjugated
NCBI	100861194
UniProt	Q29477
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Val25~Glu352 (NP_001272477.1)

Application Notes

Lactotransferrin (LTF), also known as actoferrin (LF) , is a multifunctional protein of the transferrin family. Lactoferrin belongs to the innate immune system. Apart from its main biological function, namely binding and transport of iron ions, lactoferrin also has antibacterial, antiviral, antiparasitic, catalytic, anti-cancer, and anti-allergic functions and properties. LTF is widely represented in various secretory fluids, such as milk, saliva, tears, and nasal secretions. It also present in secondary granules of PMN and is secreted by some acinar cells. Besides, Clusterin (CLU) has been identified as an interactor of LTF, thus a binding ELISA assay was conducted to detect the interaction of recombinant goat LTF and recombinant goat CLU. Briefly, LTF were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to CLU-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-LTF pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of LTF and CLU was in a dose dependent manner.