

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human JAK2 protein, His tag, Unconjugated GTX00152-PRO

Article Name	Human JAK2 protein, His tag, Unconjugated
Biozol Catalog Number	GTX00152-PRO
Supplier Catalog Number	GTX00152-pro
Alternative Catalog Number	GTX00152-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	3717
UniProt	O60674
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Asn508~Ala800 (NP_001309123.1)

Application Notes	JAK2 (Tyrosine-protein kinase JAK2) is a tyrosine kinase involved in various processes such as cell growth, development, differentiation or histone modifications. JAK2 is considered to associate with some type I receptors, including EPOR (Erythropoietin receptor), therefore participates in cellular signal transduction. Thus a binding ELISA assay was conducted to detect the interaction of recombinant human JAK2 and recombinant human EPOR. Briefly, JAK2 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl JAK2 were then transferred to EPOR-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-JAK2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of JAK2 and EPOR was in a dose dependent manner.
-------------------	--