

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human ARSB protein, His tag, Unconjugated GTX00161-PRO

Article Name	Human ARSB protein, His tag, Unconjugated
Biozol Catalog Number	GTX00161-PRO
Supplier Catalog Number	GTX00161-pro
Alternative Catalog Number	GTX00161-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	411
UniProt	P15848
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	HEK293 cells
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Leu81~Met533 (NP_000037.2)

Application Notes

Arylsulfatase B (ARSB) is a lysosomal enzyme of the sulfatase family. ARSB hydrolyzes sulfate groups of N-Acetyl-D-galactosamine, chondroitin sulfate, and dermatan sulfate. The protein is targeted to the lysosome. Besides, Plasminogen Activator, Urokinase Receptor (uPAR) has been identified as an interactor of ARSB, thus a binding ELISA assay was conducted to detect the interaction of recombinant human ARSB and recombinant human uPAR. Briefly, ARSB were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to uPAR-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-ARSB pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of ARSB and uPAR was in a dose dependent manner.