

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Cytokeratin 18 Antibody Cocktail, Clone: [Clone Cocktail], Mouse, Monoclonal NSJ-V2676-20UG**

Article Name	Cytokeratin 18 Antibody Cocktail, Clone: [Clone Cocktail], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2676-20UG
Supplier Catalog Number	V2676-20UG
Alternative Catalog Number	NSJ-V2676-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IHC-P, WB
Species Reactivity	Human
Immunogen	Recombinant full-length human KRT18 protein was used as the immunogen for the Cytokeratin 18 antibody cocktail.
Product Description	This mAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t...
Clonality	Monoclonal
Clone Designation	[Clone Cocktail]
UniProt	<a href="#">P05783</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Flow Cytometry: 1-2ug/million cells in 0.1ml,Western blot: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1),Prediluted IHC only format: incubate for 30 min at RT (2)
Application Notes	Optimal dilution of the Cytokeratin 18 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.