

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TYR Antibody Cocktail (Tyrosinase), Clone: [T311 + OCA1/812], Mouse, Monoclonal NSJ-V2915SAF-100UG

Article Name	TYR Antibody Cocktail (Tyrosinase), Clone: [T311 + OCA1/812], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2915SAF-100UG
Supplier Catalog Number	V2915SAF-100UG
Alternative Catalog Number	NSJ-V2915SAF-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for the TYR antibody cocktail.
Product Description	Recognizes a cluster of proteins between 70-80kDa, identified as Tyrosinase/TYR. Occasionally a minor band at 55kDa is also detected. This mAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is a copper-co...
Clonality	Monoclonal
Clone Designation	[T311 + OCA1/812]
UniProt	P14679
Purity	Protein G affinity chromatography

Form	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	Optimal dilution of the TYR antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.