

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

VCAM1 Antibody, Clone: [VCAM1/843], Mouse, Monoclonal NSJ-V2923-20UG

Article Name	VCAM1 Antibody, Clone: [VCAM1/843], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2923-20UG
Supplier Catalog Number	V2923-20UG
Alternative Catalog Number	NSJ-V2923-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for the VCAM1 antibody.
Product Description	Recognizes a protein of 110kDa, identified as CD106 (also known as vascular cell adhesion molecule-1 (VCAM-1), L1CAM and INCAM-100). It is a member of the Ig superfamily of adhesion molecules and is expressed at high levels on cytokine stimulated vas...
Clonality	Monoclonal
Clone Designation	[VCAM1/843]
UniProt	P19320
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Application Notes	Optimal dilution of the VCAM1 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.