

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### WT1 Antibody Cocktail, Clone: [WT1/857 + 6F-H2], Mouse, Monoclonal NSJ-V2933IHC-7ML

Article Name	WT1 Antibody Cocktail, Clone: [WT1/857 + 6F-H2], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2933IHC-7ML
Supplier Catalog Number	V2933IHC-7ML
Alternative Catalog Number	NSJ-V2933IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant full length human protein (WT1/857) and amino acids 1-181 of human Wilms Tumor (6F-H2) were used as the immunogen for the WT1 antibody cocktail.
Product Description	Recognizes a 47-55kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic ...
Clonality	Monoclonal
Clone Designation	[WT1/857 + 6F-H2]
UniProt	<a href="#">P19544</a>
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the WT1 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.