

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### WT1 Antibody Cocktail, Clone: [WT1/857 + 6F-H2], Mouse, Monoclonal NSJ-V2933SAF-100UG

Article Name	WT1 Antibody Cocktail, Clone: [WT1/857 + 6F-H2], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2933SAF-100UG
Supplier Catalog Number	V2933SAF-100UG
Alternative Catalog Number	NSJ-V2933SAF-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant full length human protein (WT1/857) and amino acids 1-181 of human Wilms Tumor (6F-H2) were used as the immunogen for the WT1 antibody cocktail.
Product Description	Recognizes a 47-55kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic ...
Clonality	Monoclonal
Clone Designation	[WT1/857 + 6F-H2]
UniProt	<a href="#">P19544</a>
Purity	Protein G affinity chromatography

Form	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	Optimal dilution of the WT1 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.