

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

MYADM Antibody, Clone: [MYADM/971], Mouse, Monoclonal NSJ-V2953IHC-7ML

Article Name	MYADM Antibody, Clone: [MYADM/971], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2953IHC-7ML
Supplier Catalog Number	V2953IHC-7ML
Alternative Catalog Number	NSJ-V2953IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for the MYADM antibody.
Product Description	Recognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellul...
Clonality	Monoclonal
Clone Designation	[MYADM/971]
UniProt	Q96S97
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the MYADM antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.