

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Myeloid Antibody / MYADM, Clone: [MYADM/972], Mouse, Monoclonal NSJ-V2954-100UG

Article Name	Myeloid Antibody / MYADM, Clone: [MYADM/972], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2954-100UG
Supplier Catalog Number	V2954-100UG
Alternative Catalog Number	NSJ-V2954-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IF, IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for the Myeloid antibody.
Product Description	Recognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellul...
Clonality	Monoclonal
Clone Designation	[MYADM/972]
UniProt	<a href="#">Q96S97</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 0.5-1ug/10 <sup>6</sup> cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Application Notes	Optimal dilution of the Myeloid antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.