

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Napsin A Antibody, Clone: [NAPSA/1239], Mouse, Monoclonal NSJ-V2996IHC-7ML

Article Name	Napsin A Antibody, Clone: [NAPSA/1239], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2996IHC-7ML
Supplier Catalog Number	V2996IHC-7ML
Alternative Catalog Number	NSJ-V2996IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	A recombinant fragment from the human protein (amino acids 189-299) was used as the immunogen for the Napsin A antibody.
Product Description	Napsin-A is a protein that in humans is encoded by the NAPSA gene. The activation peptides of aspartic proteinases plays role as inhibitors of the active site. These peptide segments, or pro-parts, are deemed important for correct folding, targeting,...
Clonality	Monoclonal
Clone Designation	[NAPSA/1239]
UniProt	<a href="#">O96009</a>
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the Napsin A antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.