

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### HCAM Antibody, Clone: [HCAM/1097], Mouse, Monoclonal NSJ-V3013-20UG

Article Name	HCAM Antibody, Clone: [HCAM/1097], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V3013-20UG
Supplier Catalog Number	V3013-20UG
Alternative Catalog Number	NSJ-V3013-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human, Mouse
Immunogen	Recombinant human protein was used as the immunogen for the HCAM antibody.
Product Description	Recognizes a cell surface glycoprotein of 80-95kDa (CD44/HCAM) on lymphocytes, monocytes, and granulocytes. The CD44 family of glycoproteins exists in a number of variant isoforms, the most common being the standard 85-95kDa or hematopoietic variant ...
Clonality	Monoclonal
Clone Designation	[HCAM/1097]
UniProt	<a href="#">P16070</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	Optimal dilution of the HCAM antibody should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.