

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

HCAM Antibody, Clone: [HCAM/1097], Mouse, Monoclonal NSJ-V3013IHC-7ML

Article Name	HCAM Antibody, Clone: [HCAM/1097], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V3013IHC-7ML
Supplier Catalog Number	V3013IHC-7ML
Alternative Catalog Number	NSJ-V3013IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human, Mouse
Immunogen	Recombinant human protein was used as the immunogen for the HCAM antibody.
Product Description	Recognizes a cell surface glycoprotein of 80-95kDa (CD44/HCAM) on lymphocytes, monocytes, and granulocytes. The CD44 family of glycoproteins exists in a number of variant isoforms, the most common being the standard 85-95kDa or hematopoietic variant ...
Clonality	Monoclonal
Clone Designation	[HCAM/1097]
UniProt	P16070
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the HCAM antibody should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.