

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### LAMP-3 Antibody / CD63, Clone: [LAMP3/968], Mouse, Monoclonal NSJ-V3029-100UG

Article Name	LAMP-3 Antibody / CD63, Clone: [LAMP3/968], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V3029-100UG
Supplier Catalog Number	V3029-100UG
Alternative Catalog Number	NSJ-V3029-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IHC-P
Species Reactivity	Human
Immunogen	Recombinant human full-length protein was used as the immunogen for the CD63 / LAMP-3 antibody.
Product Description	This mAb recognizes protein of 26kDa-60kDa, which is identified as CD63/LAMP-3. Its epitope is different from that of mAb LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cel...
Clonality	Monoclonal
Clone Designation	[LAMP3/968]
UniProt	<a href="#">P08962</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 1-2ug/10 <sup>6</sup> cells, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	Optimal dilution of the LAMP-3 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.