

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### LAMP-3 Antibody / CD63, Clone: [LAMP3/968], Mouse, Monoclonal NSJ-V3029IHC-7ML

Article Name	LAMP-3 Antibody / CD63, Clone: [LAMP3/968], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V3029IHC-7ML
Supplier Catalog Number	V3029IHC-7ML
Alternative Catalog Number	NSJ-V3029IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human full-length protein was used as the immunogen for the CD63 / LAMP-3 antibody.
Product Description	This mAb recognizes protein of 26kDa-60kDa, which is identified as CD63/LAMP-3. Its epitope is different from that of mAb LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cel...
Clonality	Monoclonal
Clone Designation	[LAMP3/968]
UniProt	<a href="#">P08962</a>
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the LAMP-3 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.