

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD63 Antibody Cocktail, Clone: [NKI/C3 + LAMP3/968], Mouse, Monoclonal NSJ-V3030IHC-7ML

Article Name	CD63 Antibody Cocktail, Clone: [NKI/C3 + LAMP3/968], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V3030IHC-7ML
Supplier Catalog Number	V3030IHC-7ML
Alternative Catalog Number	NSJ-V3030IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	The smooth plasma membrane fraction of MeWo cells (NKI/C3) and recombinant human full-length protein (LAMP3/968) were used as the immunogen for the CD63 antibody cocktail.
Product Description	This mAb recognizes protein of 26kDa-60kDa, which is identified as CD63 / LAMP-3. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes ...
Clonality	Monoclonal
Clone Designation	[NKI/C3 + LAMP3/968]
UniProt	P08962
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the CD63 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.