

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD79a Antibody, Clone: [IGA/515], Mouse, Monoclonal NSJ-V3039-100UG

| | |
|----------------------------|---|
| Article Name | CD79a Antibody, Clone: [IGA/515], Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V3039-100UG |
| Supplier Catalog Number | V3039-100UG |
| Alternative Catalog Number | NSJ-V3039-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | IF, IHC-P |
| Species Reactivity | Human |
| Immunogen | A soluble form of the extracellular IgSF domain from the human protein was used as the immunogen for the CD79a antibody. |
| Product Description | A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B... |
| Clonality | Monoclonal |
| Clone Designation | [IGA/515] |
| UniProt | P11912 |
| Purity | Protein G affinity chromatography |

| | |
|--------------------|--|
| Form | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Dilute | Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT (1) |
| Application Notes | Optimal dilution of the CD79a antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |