

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cdc20 Antibody, Clone: [CDC20/1102], Mouse, Monoclonal NSJ-V3041-100UG

Article Name	Cdc20 Antibody, Clone: [CDC20/1102], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V3041-100UG
Supplier Catalog Number	V3041-100UG
Alternative Catalog Number	NSJ-V3041-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for the Cdc20 antibody.
Product Description	Cyclins, regulatory subunits, which associate with kinases, control many of the important steps in cell cycle progression. The Cdc2 protein kinase (p34Cdc2) exhibits protein kinase activity in vitro and exists in a complex with both cyclin B and a pr...
Clonality	Monoclonal
Clone Designation	[CDC20/1102]
UniProt	Q12834
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 2-4ug/ml for 30 min at RT
Application Notes	Optimal dilution of the Cdc20 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.