

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Kappa Light Chain Antibody / Rabbit Monoclonal, Clone: [KC3L-3R] NSJ-V3713SAF-100UG

| | |
|----------------------------|---|
| Article Name | Recombinant Kappa Light Chain Antibody / Rabbit Monoclonal, Clone: [KC3L-3R] |
| Biozol Catalog Number | NSJ-V3713SAF-100UG |
| Supplier Catalog Number | V3713SAF-100UG |
| Alternative Catalog Number | NSJ-V3713SAF-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC-P |
| Species Reactivity | Human |
| Immunogen | Recombinant human Ig kappa chain was used as the immunogen for this recombinant Kappa light chain antibody. |
| Product Description | This mAb is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. In mammals, the two light chains in an antibody are always identical, with only one type of light chain, ... |
| Clonality | Monoclonal |
| Clone Designation | [KC3L-3R] |
| UniProt | P01601 |
| Purity | Protein A affinity chromatography |

| | |
|--------------------|--|
| Form | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Dilute | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT, Prediluted IHC only format: incubate for 30 min at RT (1) |
| Application Notes | The optimal dilution of the recombinant Kappa light chain antibody for each application should be determined by the researcher. 1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |