

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-CD56 Antibody, Clone: [SPM489], Mouse, Monoclonal NSJ-V9055SAF-100UG

Article Name	Anti-CD56 Antibody, Clone: [SPM489], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9055SAF-100UG
Supplier Catalog Number	V9055SAF-100UG
Alternative Catalog Number	NSJ-V9055SAF-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	A membrane preparation of small cell lung carcinoma was used as the immunogen for this anti-CD56 antibody. This mAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM.
Product Description	Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135kDa isoform is the basic molecule, which is glycosylated or sialylated to produce the mature species. Anti-C...
Clonality	Monoclonal
Clone Designation	[SPM489]
UniProt	P13591
Purity	Protein G affinity chromatography

Form	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	The optimal dilution of the anti-CD56 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.