

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Ornithine Decarboxylase Antibody, Clone: [SPM565], Mouse, Monoclonal NSJ-V9062-20UG

Article Name	Ornithine Decarboxylase Antibody, Clone: [SPM565], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9062-20UG
Supplier Catalog Number	V9062-20UG
Alternative Catalog Number	NSJ-V9062-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for this ODC-1 antibody.
Product Description	Recognizes a 53kDa protein, identified as the Ornithine Decarboxylase (ODC-1). ODC is the initial and rate-limiting enzyme in the biosynthetic pathway of polyamines and is involved in the conversion of ornithine to putrescine. The biological activity...
Clonality	Monoclonal
Clone Designation	[SPM565]
UniProt	P11926
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	The optimal dilution of the Ornithine Decarboxylase antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.