

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-PCNA Antibody, Clone: [SPM350], Mouse, Monoclonal NSJ-V9064-100UG

Article Name	Anti-PCNA Antibody, Clone: [SPM350], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9064-100UG
Supplier Catalog Number	V9064-100UG
Alternative Catalog Number	NSJ-V9064-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P, WB
Species Reactivity	Human
Immunogen	Rat PCNA protein was used as the immunogen for this anti-PCNA antibody.
Product Description	Recognizes a non-histone protein of 36kDa, which is identified as proliferating cell nuclear antigen (PCNA). It is also known as cyclin or polymerase delta auxiliary protein. Elevated expression of PCNA/cyclin has been shown in the nucleus during lat...
Clonality	Monoclonal
Clone Designation	[SPM350]
UniProt	P12004
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Western blot: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	The optimal dilution of the anti-PCNA antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.