

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-Bcl-X Antibody, Clone: [SPM165], Mouse, Monoclonal NSJ-V9075IHC-7ML

Article Name	Anti-Bcl-X Antibody, Clone: [SPM165], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9075IHC-7ML
Supplier Catalog Number	V9075IHC-7ML
Alternative Catalog Number	NSJ-V9075IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Amino acids 3-14 (QSNRELVVDFLS) from the human protein were used as the immunogen for this anti-Bcl-X antibody.
Product Description	Recognizes a protein of 27kDa, identified as the Bcl-X protein. This mAb shows no cross-reaction with Bcl-2 or Bax protein. Bcl-X has two isoforms, Bcl-XL (long), a 241 amino acid protein which suppresses cell death. And Bcl-XS (short), a 178 amino a...
Clonality	Monoclonal
Clone Designation	[SPM165]
UniProt	<a href="#">Q07817</a>
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	The optimal dilution of the anti-Bcl-X antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.