

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Rabbit IgM (μ)-HRPO, MinX none DNA-SEC-182606

Article Name	Goat IgG anti-Rabbit IgM (μ)-HRPO, MinX none
Biozol Catalog Number	DNA-SEC-182606
Supplier Catalog Number	SEC-182606
Alternative Catalog Number	DNA-SEC-182606
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Rabbit
Immunogen	Anti-Rabbit IgM (mu) was produced by repeated immunization with rabbit IgM mu heavy chain in goat.
Conjugation	HRPO
Format	IgG
Target Specificity	IgM (μ)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Rabbit IgM (mu) peroxidase conjugated antibody generated in goat detects specifically rabbit IgM (mu). This secondary peroxidase conjugated antibody anti-Rabbit is ideal for investigators who routinely perform titration assays, western-blot, imm...
Clonality	Polyclonal

Concentration	10.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by cross adsorption to remove any unwanted reactivities and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Rabbit IgM and Rabbit Serum. No reaction was observed against Rabbit IgG.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Rabbit
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Application Notes	Rabbit IgM (mu) peroxidase conjugated antibody has been assayed against 1.0 ug of Rabbit IgM in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:40,000 of the reconstitution concentration is suggested for this product.