

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit F(ab)2 anti-Dog IgG (F(ab)2)-HRPO, MinX none DNA-SEC-182665

Article Name	Rabbit F(ab)2 anti-Dog IgG (F(ab)2)-HRPO, MinX none
Biozol Catalog Number	DNA-SEC-182665
Supplier Catalog Number	SEC-182665
Alternative Catalog Number	DNA-SEC-182665
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Canine
Immunogen	Dog IgG F(ab)2 fragment
Conjugation	HRPO
Format	F(ab')2
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	F(ab)2 Anti-Dog IgG F(ab)2 Peroxidase Antibody generated in rabbit detects Dog F(ab)2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secr...
Clonality	Polyclonal

Concentration	10.0 mg/mL
Isotype	Ig
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product is a F(ab') ₂ fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Dog IgG, Dog IgG F(ab') ₂ and Dog Serum. No reaction was observed against Dog IgG F(c), anti-Rabbit IgG F(c) or anti-Pepsin.
Form	Lyophilized
Formula	10 mM NaPO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Gentamicin
Target	Dog
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Application Notes	This product has been assayed against 1.0 ug of Dog IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product.