

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Chicken IgG (H+L)-unconj., MinX Bo,Go,Gp,Hm,Ho,Hu,Ms,Rb,Rt,Sh DNA-SEC-182775

Article Name	Goat IgG anti-Chicken IgG (H+L)-unconj., MinX Bo,Go,Gp,Hm,Ho,Hu,Ms,Rb,Rt,Sh
Biozol Catalog Number	DNA-SEC-182775
Supplier Catalog Number	SEC-182775
Alternative Catalog Number	DNA-SEC-182775
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Gallus
Immunogen	Chicken IgG whole molecule
Conjugation	Unconjugated
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Goat,Guinea pig,Hamster (all),Equine,Human,Mouse,Rabbit,Rat,Sheep
Product Description	Anti-Chicken IgG Antibody generated in goat detects chicken IgY. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fu...

Clonality	Polyclonal
Concentration	1.1 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by conjugation to fluorochrome and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Chicken IgG and Chicken Serum. No reaction was observed against Bovine, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit, Rat and Sheep Serum Proteins. This antibody will react with heavy chains of Chicken IgG and with light chains of most Chicken immunoglobulins.
Form	Liquid (sterile filtered)
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Chicken
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:600,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Anti-Chicken IgG antibody has been tested by ELISA and western blot and is suitable for immunohistochemistry, as well as other assays requiring lot-to-lot consistency.