

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Chicken IgG (H+L)-FITC, MinX Bo,Go,Gp,Hm,Ho,Hu,Ms,Rb,Rt,Sh DNA-SEC-182778

Article Name	Goat IgG anti-Chicken IgG (H+L)-FITC, MinX Bo,Go,Gp,Hm,Ho,Hu,Ms,Rb,Rt,Sh
Biozol Catalog Number	DNA-SEC-182778
Supplier Catalog Number	SEC-182778
Alternative Catalog Number	DNA-SEC-182778
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	FLISA,FACS,IF
Species Reactivity	Gallus
Immunogen	Chicken IgG whole molecule
Conjugation	FITC
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Goat,Guinea pig,Hamster (all),Equine,Human,Mouse,Rabbit,Rat,Sheep
Product Description	Anti-Chicken IgG Fluorescein Antibody generated in goat detects chicken IgY. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, a...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by conjugation to fluorochrome and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-fluorescein, anti-Goat Serum, Chicken IgG and Chicken Serum. No reaction was observed against Bovine, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit, Rat and Sheep Serum Proteins. This antibody will react with heavy chains of Chicken IgG and with light chains of most Chicken immunoglobulins.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Chicken
Antibody Type	Secondary Antibody
Application Dilute	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 2.8, IF Microscopy Dilution: 1:1,000 - 1:5,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.