

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Horse IgG (F(ab)2)-Biotin, MinX none DNA-SEC-182971

Article Name	Goat IgG anti-Horse IgG (F(ab)2)-Biotin, MinX none
Biozol Catalog Number	DNA-SEC-182971
Supplier Catalog Number	SEC-182971
Alternative Catalog Number	DNA-SEC-182971
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Equine
Immunogen	Horse IgG F(ab)2 fragment
Conjugation	Biotin
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Horse IgG F(ab)2 Biotin Antibody generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. F(ab)2 molecules lack the Fc por...
Clonality	Polyclonal

Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Horse IgG, Horse IgG F(ab') ₂ and Horse Serum. No reaction was observed against Horse IgG F(c).
Form	Lyophilized
Formula	20 mM K ₃ PO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% NaN ₃
Target	Horse
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	This product has been assayed against 1.0 ug of Horse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.