

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Rat IgG (H+L)-FITC, MinX none DNA-SEC-183496

Article Name	Rabbit IgG anti-Rat IgG (H+L)-FITC, MinX none
Biozol Catalog Number	DNA-SEC-183496
Supplier Catalog Number	SEC-183496
Alternative Catalog Number	DNA-SEC-183496
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	FLISA,FACS,IF
Species Reactivity	Rat
Immunogen	Rat IgG whole molecule
Conjugation	FITC
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Rat IgG (H&L) fluorescein Antibody generated in rabbit detects reactivity to Rat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruse...
Clonality	Polyclonal

Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	Anti-Rat IgG (H&L) fluorescein Antibody generated in rabbit detects reactivity to Rat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both the Heavy and Light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. This Anti-Rat IgG antibody is conjugated to Fluorescein.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Rat
Antibody Type	Secondary Antibody
Application Dilute	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 3.1, IF Microscopy Dilution: 1:1,000 - 1:5,000
Application Notes	Secondary antibody reagents are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.