

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Sheep IgG (Fc)-Alk. Phos., MinX Hu DNA-SEC-183559

Article Name	Rabbit IgG anti-Sheep IgG (Fc)-Alk. Phos., MinX Hu
Biozol Catalog Number	DNA-SEC-183559
Supplier Catalog Number	SEC-183559
Alternative Catalog Number	DNA-SEC-183559
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Sheep
Immunogen	Sheep IgG F(c) fragment
Conjugation	Alk. Phos.
Format	IgG
Target Specificity	IgG (Fc)
Cross-Adsorption (MinX)	Human
Product Description	Anti-Sheep IgG F(c) generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of sheep IgG and o...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Sheep IgG, Sheep IgG F(c) and Sheep Serum. No reaction was observed against Sheep IgG F(ab) or Human Serum Proteins..
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target	Sheep
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Application Notes	This product has been assayed against 1.0 ug of Sheep IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:14,000 of the reconstitution concentration is suggested for this product.