

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Swine IgG (F(ab)2)-Alk. Phos., MinX none DNA-SEC-183590

Article Name	Rabbit IgG anti-Swine IgG (F(ab)2)-Alk. Phos., MinX none
Biozol Catalog Number	DNA-SEC-183590
Supplier Catalog Number	SEC-183590
Alternative Catalog Number	DNA-SEC-183590
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Porcine
Immunogen	Swine IgG F(ab)2 fragment
Conjugation	Alk. Phos.
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Swine IgG F(ab)2 Antibody generated in rabbit recognizes the dimeric Fab portion of the Swine IgG molecule. Swine IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlle...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Swine IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Swine IgG, Swine IgG F(ab') ₂ and Swine Serum. No reaction was observed against Swine IgG F(c).
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl, 150 mM NaCl, 1 mM MgCl ₂ , 0.1 mM ZnCl ₂ , 50% (v/v) Glycerol, pH 8.0, sterile filtered, 0.01% NaN ₃
Target	Swine
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:1,000 - 1:3,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,000
Application Notes	Anti-Swine IgG F(ab) ₂ alkaline phosphatase conjugated antibody has been tested by ELISA and is suitable for use in ELISA, immunohistochemistry, and western blot. Specific conditions for reactivity should be optimized by the end user.